

Healthy eating

eating while you are pregnant

choosing food
to keep you
and your baby
healthy




FOOD
STANDARDS
AGENCY

During pregnancy you need to make sure that your diet is providing enough energy and nutrients for your baby to grow and develop, and for your body to cope with the changes taking place. This booklet contains practical advice to help you make some healthy food choices while you're pregnant.

Q What should I be eating?

A It's important to try to eat a variety of foods including:

- plenty of fruit and vegetables (fresh, frozen, tinned, dried or a glass of juice) – aim for at least five portions of a variety of fruit and vegetables a day
- plenty of starchy foods, such as bread, pasta, rice and potatoes
- some protein, such as lean meat and chicken, fish (aim for at least two servings of fish a week, including one of oily fish), eggs and pulses (such as beans, peas and lentils). These are also good sources of iron (see page 2)
- plenty of fibre, found in wholegrain bread, wholegrain cereals, pasta, rice, pulses, and fruit and vegetables – this helps prevent constipation
- some dairy foods such as milk, cheese and yoghurt, which contain calcium. These are also good sources of protein



Q Do I need extra iron?

- A Pregnant women can become short of iron, so make sure you choose plenty of iron-rich foods. Try to have some food or drink containing vitamin C, such as a glass of fruit juice, at the same time as an iron-rich meal because this might help your body absorb the iron.



Avoid drinking tea or coffee with iron-rich foods because these could make it harder for the body to absorb iron.

Good sources of iron

- red meat
- pulses
- bread
- green leafy vegetables
- fortified breakfast cereals
- dried fruit

Although liver also contains lots of iron, you should avoid eating it while you're pregnant. (See **Is there any food I should avoid?** on page 4.)

If the iron level in your blood becomes low, your GP or midwife may advise you to take iron supplements.



Q Do I need to take any vitamin supplements?

You should take a 400 microgram (mcg) folic acid supplement each day until the 12th week of your pregnancy. Ideally, you should have started to take these supplements before you were pregnant.

You should also include foods containing folic acid in your diet, such as green leafy vegetables, fortified breakfast cereals and brown rice.

Folic acid has been shown to reduce the risk of neural tube defects such as spina bifida. If you want to get your folic acid from a multivitamin tablet, make sure it contains 400 mcg and does not contain vitamin A (or retinol). Remember that if you take more than one multivitamin tablet, you could overdose on the other vitamins they contain. (See **Is there any food I should avoid?** on page 4.)

If you have already had a pregnancy affected by a neural tube defect, consult your GP for advice.

You also need to take supplements containing 10 mcg of vitamin D each day.

If you or your family get Income Support, Income-based Jobseeker's Allowance or Child Tax Credit, you may be able to get free vitamin supplements through Healthy Start. Check with your midwife.

Oily fish, eggs and some fat spreads provide vitamin D in our diet but most of our vitamin D comes from the action of summer sunlight on our skin. But remember, if you're out in the sun, take care not to burn.

If you have dark skin or if you always cover up all your skin when you're outside, you may be particularly short of vitamin D. Ask your GP for more information.

Q Is there any food I should avoid?

A There are certain foods you shouldn't eat while you're pregnant because they can contain high levels of listeria, a germ that can cause miscarriage, stillbirth or severe illness in a newborn baby. Remember to avoid the following:

- cheeses, such as Camembert, Brie or chevre (a type of goats' cheese), or others that have a similar rind. You should also avoid blue cheese. You don't need to avoid hard cheeses (such as Parmesan and Cheddar), cottage cheese and processed cheese
- pâté (any type, including vegetable)
- uncooked or undercooked ready meals. Make sure you heat ready meals until they're piping hot all the way through

You should also remember to:

- Avoid eating raw eggs and food containing raw or partially cooked eggs. Only eat eggs that have been cooked until both the white and yolk are solid. This is to avoid the risk of salmonella, which causes a type of food poisoning.
- Always wash your hands after handling raw meat, and keep raw foods separate from ready-to-eat foods. This is to avoid food poisoning germs, such as salmonella, campylobacter and *E. coli* O157.
- Make sure you only eat meat that has been well cooked. Take particular care with sausages and minced meat.
- Always wear gloves when you're gardening or changing cat litter, and wash your hands afterwards. This is to avoid toxoplasmosis, an infection caused by a parasite found in meat, cat faeces and soil. Toxoplasmosis can be harmful to unborn babies.



- Make sure you don't have too much vitamin A. This means you should avoid taking supplements containing vitamin A and fish liver oil supplements, and avoid eating liver and liver products such as pâté. You need some vitamin A, but too much could harm your baby. Ask your GP or midwife if you would like more information.
- Avoid eating shark, swordfish and marlin. And limit the amount of tuna you eat to no more than two steaks a week (weighing about 140g cooked or 170g raw) or four medium-size cans a week (about 140g drained weight per can). This is because of the levels of mercury in these fish. At high levels, mercury can harm a baby's developing nervous system.
- Have no more than two portions of oily fish a week. Oily fish includes fresh tuna (not canned tuna), mackerel, sardines and trout.
- Choose cooked shellfish rather than raw. This is because raw shellfish might be contaminated with harmful bacteria and viruses that could cause food poisoning. And food poisoning can be particularly unpleasant during pregnancy.

It's also a good idea to:

- Cut down on foods such as cakes and biscuits, because these are high in fat and sugar. This can also help you to avoid putting on too much weight during pregnancy.

Some healthier light meals and snacks

- sandwiches or pitta bread filled with chicken, cottage cheese or lean ham
- low-fat yoghurts
- vegetable and bean soups
- fruit – fresh, canned in juice or dried fruit such as raisins



Q Do I need to cut out caffeine?

A You should limit the amount of caffeine you have each day, but you don't need to cut it out completely. Caffeine occurs naturally in some food and hot drinks, such as coffee, tea and chocolate, and it's also added to some soft drinks and 'energy' drinks.

It's important to have no more than 300 mg of caffeine a day. This is because high levels of caffeine can lead to low birth weight, or even miscarriage.

Each of these contains roughly 300 mg:

- 3 mugs of instant coffee (100 mg each)
- 4 cups of instant coffee (75 mg each)
- 3 cups of brewed coffee (100 mg each)
- 6 cups of tea (50 mg each)
- 8 cans of cola (up to 40 mg each)
- 4 cans of 'energy' drink (up to 80 mg each)
- 8 bars of chocolate, weighing 50 g each (up to 50 mg each)



So, if you eat a bar of chocolate and drink 3 cups of tea, a can of cola and a cup of instant coffee in a day, you'll have reached the 300 mg maximum amount of caffeine.

Remember that caffeine is also found in certain cold and flu remedies. Always check with your GP or another health professional before taking any medication or over-the-counter remedies.

Q Do I need to cut out alcohol?

- A It's best to stop drinking altogether when you're pregnant. But if you do drink, limit yourself to no more than 1 or 2 units of alcohol, once or twice a week and avoid getting drunk. A unit is half a pint of ordinary-strength beer, lager or cider, or a single 25 ml measure of spirits. A glass of wine is about 2 units and alcopops are about 1.5 units.

Q How much weight should I expect to put on?

- A Weight gain varies a great deal and depends on what you weighed before you became pregnant. But most women put on 10 to 12.5 kg (22 to 28 lb) over the whole of their pregnancy.

If you gain too much weight, this can affect your health and increase your blood pressure. But equally, it's important that you don't try to diet.

Cutting down on fatty and sugary food and drink may help you to avoid gaining excessive weight during pregnancy.



Q Where can I get more information?

- A You might find it useful to read *The pregnancy book*, which is published by health departments in England, Wales and Northern Ireland, and is available free to first-time parents. Health Scotland produces a book called *Ready Steady Baby!*, which is free to all pregnant women in Scotland.

If you haven't already got a copy of one of these, speak to your midwife, health visitor or GP, or contact your local health promotion unit (or department in Scotland).

You can find out more about food at eatwell.gov.uk





eatwell.gov.uk/pregnancy

For more information and advice about food,
visit the Food Standards Agency's websites:

eatwell.gov.uk

food.gov.uk

salt.gov.uk

Food Standards Agency Publications

To order copies of this or other publications produced
by the Agency, contact Food Standards Agency Publications:

tel **0845 606 0667**
minicom **0845 606 0678**
fax **020 8867 3225**
email **foodstandards@ecgroup.co.uk**

Published by the Food Standards Agency 2002.
Reprinted with amendments March 2008

Printed by Oakfield Press Ltd 100K

© Crown copyright 2002

Printed in England on paper comprising a minimum of 75% recycled fibre
FSA/0451/0308



When you have finished with
this publication please recycle it