

Development of accurate predictive models for the assessment of the survival of *Campylobacter jejuni* and *Campylobacter coli*.

Area of research interest: [Foodborne pathogens](#)

Study duration: 2011-04-01

Planned completion: 1 January 2015

Project code: FS241040

Conducted by: University of Liverpool

Background

Campylobacter is the leading cause of foodborne disease in the UK and tackling this problem is a key priority for the Agency. *Campylobacter* is present on a significant proportion of raw poultry meat entering kitchens and poor hygiene can lead to cross-contamination of surfaces, utensils, hands and ready-to-eat foods.

Predictive models for growth and survival of foodborne pathogens are publically available and provide industry, government scientists and researchers with tools to predict pathogen behaviour in response to changes in food formulation and storage conditions. Models are available for most foodborne pathogens; and at the time of commissioning this project there was a paucity of predictive models for *Campylobacter*. Since then more data has been produced, which is available in ComBase.

Given the paucity of data at the time, research was needed to develop robust predictive modelling tools for *Campylobacter* growth and survival in food, particularly thermal death.

Research Approach

The project aimed to develop predictive models for the growth and survival of *Campylobacter* in food that could be used by the food industry to address this hazard in food production and inform instructions for caterers/consumers regarding appropriate handling, cooking and storage of foods which may act as a vehicle for *Campylobacter*.

A systematic review of published literature on the survival of *Campylobacter* in foods and food-related environments was carried out to identify data gaps in this area.

The study also examined the heat resistance of *Campylobacter* at the population level and used isolates from food and humans. It also involved an investigation into how *Campylobacter* interacts with food and used highly sensitive recovery and enumeration techniques. A generic modelling approach was used to develop predictive models for growth and survival across all of the different treatment, food states and environments investigated, using the data collected from these

experiments.

There has been an increase in the use of sous vide cooking, which involves vacuum packing food and cooking in a water bath at lower temperatures over a prolonged period of time. Currently there is a gap in the data on the effect of cooking food at lower temperatures and on the growth and inactivation of *Campylobacter*. It is unclear whether this presents a food safety risk. The ACMSF have recommended that more research is required to look at lower cooking temperatures and reliably establish safe time/temperature combinations to reduce potential risks to consumers. As this research was looking at the thermal inactivation of *Campylobacter*, this work was extended to include lower temperatures to help fill the research gap associated with the issue of sous vide cooking.

These models will be made publically available, and in conjunction with the data available on ComBase, will help the food industry address this hazard in food production and inform instructions for caterers/consumers regarding handling, cooking and storage of foods which may act as a vehicle for *Campylobacter*.

Results

This project generated a number of results/conclusions and the main ones have been summarised below:

1. Data generated in this project show that *Campylobacter* can survive at lower pH ranges (e.g. pH 5.5) when exposed to high temperatures.
2. *Campylobacter* attachment to chicken meat enhanced its survival only at sub-lethal temperatures (e.g. 56°C), with pre-chilling having limited impact on sensitivity to heat treatment.
3. Gradual heating of chicken meat pieces inoculated with high levels externally and internally to 70°C has demonstrated survival of low numbers of *Campylobacter*, which is of potential concern.
4. Given that retail surveys have found levels of up to 105 CFU/g contamination of *Campylobacter* on carcasses, these experimental results may suggest that the current recommendation of 70°C for two minutes may be insufficient to inactivate all *Campylobacter* present in chicken meat when contaminated at such levels. However, more work is required to investigate if such sub-populations of viable cells are still able to cause infection. Further work should also be undertaken to elucidate the mechanisms involved in the survival of these residual populations and their association with chicken meat surfaces.
5. Cooking chicken at lower temperatures (50 and 52°C) in a vacuum sealed pack for three and two hours respectively was inadequate in eliminating *Campylobacter*, although it should be noted that a high inoculum was used in these studies. *Campylobacter* was still isolated from whole chicken fillets following these heat treatments, with *C. coli* also recovered at 56°C after one hour of cooking.
6. Our findings indicate that no single model describes the survival of *Campylobacter* under all circumstances, and that the selection of an appropriate model should be based on individual strain and response to the type and magnitude of challenge used during experimental simulation.

The findings of this report are interesting; however there are a number of criticisms around the methodology used, such as the use of a water bath in the heating of chicken meat not being a true reflection on actual cooking practices. In addition, the project did not produce a model that was as user-friendly as anticipated, however this was due to the fact that a one-model-fits-all approach could not be achieved as the different strains used in the experimental work produced varied responses. The suggestion that the current cooking advice of 70°C for 2 minutes may be insufficient should not be interpreted on its own and should be considered alongside numerous

other studies that provide a body of evidence to show this advice to be effective in eliminating *Campylobacter*. It is therefore unlikely that FSA will be changing its current cooking advice based on this one study. This data should be uploaded into ComBase to add to the existing data already available on *Campylobacter*, allowing for a much more user-friendly interface and a more realistic prediction.

Related media

England, Northern Ireland and Wales

PDF

[View FS241040 - Contents as PDF\(Open in a new window\)](#) (1.18 MB)

England, Northern Ireland and Wales

PDF

[View Development of accurate predictive models for the assessment of the survival of *Campylobacter jejuni* and *C* as PDF\(Open in a new window\)](#) (504.52 KB)

England, Northern Ireland and Wales

PDF

[View Chapter 1 Development and implementation of Non-linear models within R framework as PDF\(Open in a new window\)](#) (3.95 MB)

England, Northern Ireland and Wales

PDF

[View Chapter 2 Development and implementation of Non-linear models within GlnaFit framework as PDF\(Open in a new window\)](#) (2.19 MB)

England, Northern Ireland and Wales

PDF

[View Chapter 2b - Food-matrices Exterior Time-temperature Simulations: Gradual Heating 56o C as PDF\(Open in a new window\)](#) (1.36 MB)

England, Northern Ireland and Wales

PDF

[View Chapter 3 Sous-Vide time-temperature profiles as PDF\(Open in a new window\)](#) (1000.65 KB)

England, Northern Ireland and Wales

PDF

[View Appendices Description of the parameter set estimated by the asymptotic regression model as PDF\(Open in a new window\)](#) (753.06 KB)

England, Northern Ireland and Wales

PDF

[View GlnaFit A user guide as PDF\(Open in a new window\)](#) (1.01 MB)

England, Northern Ireland and Wales

PDF

[View Development of accurate predictive models for the assessment of the survival of Campylobacter jejuni and C. coli as PDF\(Open in a new window\)](#) (1.24 MB)

England, Northern Ireland and Wales

EXCEL

[View Time-temperature 56C All Data as Excel\(Open in a new window\)](#) (335.01 KB)

England, Northern Ireland and Wales

EXCEL

[View Time-temperature Simulations 60C Data as Excel\(Open in a new window\)](#) (357.21 KB)

England, Northern Ireland and Wales

EXCEL

[View Time-temperature 64C All Data as Excel\(Open in a new window\)](#) (367.02 KB)

EXCEL

[View FS 241040 Growth limit detection data as Excel\(Open in a new window\)](#) (11.19 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - CAB Media Detection Data as Excel\(Open in a new window\)](#) (136.07 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Sous Vide Whole-fillet Data 50C as Excel\(Open in a new window\)](#) (25.4 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Sous Vide Whole-fillet Data 52C as Excel\(Open in a new window\)](#) (92.97 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Sous Vide Whole-fillet Data 54C as Excel\(Open in a new window\)](#) (41.73 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Sous Vide Internal Survival Data as Excel\(Open in a new window\)](#) (47.13 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - PH Time-temperature Simulations 56C Data as Excel\(Open in a new window\)](#) (586.16 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - PH Time-temperature Simulations 60C Data as Excel\(Open in a new window\)](#)
(415.99 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - PH Time-temperature Simulations 64C Data as Excel\(Open in a new window\)](#)
(450.93 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Gradual Heating 56C as Excel\(Open in a new window\)](#) (202.83 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Gradual Heating 70C as Excel\(Open in a new window\)](#) (152.58 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Interiors Direct Heating 64C as Excel\(Open in a new window\)](#)
(98.7 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Interiors Direct Heating 68C as Excel\(Open in a new window\)](#)
(101.28 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Direct Heating 56C as Excel\(Open in a new window\)](#) (204.63 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Direct Heating 60C as Excel\(Open in a new window\)](#) (418.23 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Direct Heating 64C as Excel\(Open in a new window\)](#) (205.18 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Direct Heating 68C as Excel\(Open in a new window\)](#) (150.5 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Direct Heating 70C as Excel\(Open in a new window\)](#) (304.51 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Interiors Gradual Heating 64C as Excel\(Open in a new window\)](#) (50 KB)

England, Northern Ireland and Wales

EXCEL

[View FS241040 - Food Matrices Interiors Gradual Heating 68C as Excel\(Open in a new window\)](#) (90.02 KB)